

Comment on “Nature of the Spin Glass State”

C.M. Newman

Courant Inst. of Math. Sciences, New York Univ., New York, NY 10012

D.L. Stein

Departments of Physics and Mathematics, University of Arizona, Tucson, AZ 85721

We show that the interface excitations of Palassini-Young and Krzakala-Martin cannot yield new thermodynamic states.

75.10.Nr, 05.70.Jk, 75.50.Lk, 75.40.Mg

Recently, Palassini and Young [1] numerically studied the ground state structure of nearest-neighbor Ising spin glasses with Gaussian couplings in spatial dimensions $d = 3, 4$. Using a novel coupling-dependent bulk term in the Hamiltonian, they looked for large-scale, low-energy excitations above the ground state in fixed volume with periodic boundary conditions (b.c.’s). They interpret their results as evidence of spin configurations whose interface with the ground state (i.e., couplings satisfied in the excitation configuration but not in the ground state, or vice-versa) scales as L^{d_s} , with $d_s < d$. At the same time the excitation energy was found to scale as $L^{\theta'}$, with $\theta' = 0$; i.e., it remained finite as $L \rightarrow \infty$ [2]. This is in distinction to the similar exponent θ , which governs the energy difference under a change in coupling-independent b.c.’s, such as periodic to antiperiodic. In both $d = 3, 4$, numerical studies are consistent with $\theta > 0$ (see references in [1]). Similar results to [1] were found in $d = 3$ by Krzakala and Martin [3].

In this Comment, we show that the two properties $d_s < d$ and $\theta' = 0$ are incompatible thermodynamically; i.e., such excitations cannot give rise to new (ground or pure) states: if such interfaces exist, they cannot be pinned, but rather must deflect to infinity as $L \rightarrow \infty$. (They could also vanish altogether, as noted in [1].)

Suppose that these interfaces are pinned; this implies the existence of infinite volume states that differ by interfaces with $d_s < d$ and whose energy remains of order one as $L \rightarrow \infty$. Consider two such states α and β and the periodic b.c. metastate [4,5] for the above model with periodic b.c.’s (our argument works also for other b.c.’s chosen independently of the couplings); the metastate simply gives the probability of finding a given ground state pair (GSP) in a typical large volume.

As one moves along the interface, its energy should be larger sometimes in α and sometimes in β , so in some volumes α would be the ground state and in others β . Both α and β would therefore have positive weight in the metastate, i.e., would appear in a positive fraction of volumes. For simplicity we assumed above only a single excitation, but this argument can be extended to an

arbitrary number (even uncountably many).

But, as in [6] (Lemma 2), if the periodic b.c. metastate is supported on multiple GSP’s, these must have relative interfaces with dimension $d_s = d$. It follows that the conditions $d_s < d$ and $\theta' = 0$ are incompatible for pinned interfaces on sufficiently large length scales. (Pinned interfaces with $d_s < d$, leading to multiple ground states at zero temperature and pure states at positive temperature, would probably require $\theta' > (d - 1)/2$ [7–9], and could only be seen with coupling-dependent b.c.’s.)

It follows that, if there exist excitations above the ground state with $d_s < d$ and energy of order one independently of L , they would simply deflect to infinity as $L \rightarrow \infty$. So the possibility of their existence could be relevant in determining the excitation spectrum of spin glasses, but they would not imply the existence of new thermodynamic (ground or pure) states.

This research was supported in part by NSF Grants DMS-98-02310 (CMN) and DMS-98-02153 (DLS). The authors thank Peter Young and Matteo Palassini for useful correspondence.

-
- [1] M. Palassini and A.P. Young, Phys. Rev. Lett. **85**, 3017 (2000). Also available as cond-mat/0002134.
 - [2] There has been some dispute about these results based on interpretation of numerical data; see A.A. Middleton, cond-mat/0007375 and E. Marinari and G. Parisi, cond-mat/0007493. However, the analysis presented here is based on general thermodynamic arguments.
 - [3] F. Krzakala and O.C. Martin, Phys. Rev. Lett. **85**, 3013 (2000). Also available as cond-mat/0002055.
 - [4] C.M. Newman and D.L. Stein, Phys. Rev. Lett. **76**, 4821 (1996).
 - [5] C.M. Newman and D.L. Stein, Phys. Rev. E **55**, 5194 (1997).
 - [6] C.M. Newman and D.L. Stein, Phys. Rev. Lett. **84**, 3966 (2000).
 - [7] D.S. Fisher and D.A. Huse, Phys. Rev. B **38**, 386 (1988).
 - [8] A. van Enter, J. Stat. Phys. **60**, 275 (1990).
 - [9] C.M. Newman and D.L. Stein, Phys. Rev. B **46**, 973 (1992).